



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

is valid for any real value of n . Since in the above considerations no restrictions were put upon x , (F) holds for any value of x , real or complex, if we assume, as we may, that (f) holds for such values of the variables. Thus we may write:

$$(8) \quad f(z) = f(u + vi) = f(u) + f(vi) + 2cuv i.$$

But

$$f(u) = f(u \cdot 1) = uf(1) + u(u-1)c(1)^2,$$

$$f(vi) = f(v \cdot i) = vf(i) + v(v-1)c(i)^2;$$

hence, (8) may be written:

$$(9) \quad f(z) = f(u + vi) = uf(1) + vf(i) + c(u + vi)^2 - (u - v)c.$$

In order to determine the value of $f(1)$ we put $u = 1$, and $v = 0$ and we have from (9) $f(1) = f(1)$, which shows that the value of $f(1)$ is arbitrary. By putting $u = 0$, $v = 1$, we may show in the same way that $f(i)$ is arbitrary. Hence denoting $f(1) - c$ by α and $f(i) + c$ by β , where α and β are two arbitrary quantities, (9) may be written

$$f(u + vi) = c(u + vi)^2 + \alpha u + \beta v.$$

If $\alpha = \beta = 0$, i. e., if $f(1) = 1$ and $f(i) = -1$, and $c = 1$ we have

$$f(u + vi) = (u + vi)^2.$$

Also variously solved by T. M. SIMPSON, O. S. ADAMS, ELIJAH SWIFT, E. R. SMITH, HORACE OLSON, A. A. BENNETT, C. F. GUMMER, and J. L. WALSH.

468. Proposed by H. C. FEEMSTER, York College, Neb.

In each of the following series find the n th term and sum:

$$(a) \quad 2 + 5 + 9 + 15 + 24 + \cdots,$$

$$(b) \quad 1 + 6 + 10 + 20 + 35 + \cdots,$$

$$(c) \quad 1 + 5 + 15 + 35 + 70 + \cdots,$$

SOLUTION BY J. L. RILEY, Tahlequah, Okla.

(a) Using the method of differences we have

$$\begin{array}{ccccccc} 3 & 4 & 6 & 9 & \cdots \\ & 1 & 2 & 3 & \cdots \\ & & 1 & 1 & \cdots \\ & & & 0 & \cdots \end{array}$$

$$\begin{aligned} U_n &= 2 + 3(n-1) + \frac{(n-1)(n-2)}{2} + \frac{(n-1)(n-2)(n-3)}{3} \\ &= \frac{n^3 - 3n^2 + 20n - 6}{6}, \text{ the } n\text{th term,} \end{aligned}$$

$$\begin{aligned} S_n &= 2n + \frac{3n(n-1)}{2} + \frac{n(n-1)(n-2)}{3} + \frac{n(n-1)(n-2)(n-3)}{4} \\ &= \frac{n}{24} (n^3 - 2n^2 + 35n + 14), \text{ the sum.} \end{aligned}$$

(b) In the series $1 + 6 + 10 + 20 + 35 + \cdots$, let $U_n = A + Bn + Cn^2 + Dn^3 + En^4$. Then

$$\begin{cases} A + B + C + D + E = 1, \\ A + 2B + 4C + 8D + 16E = 6, \\ A + 3B + 9C + 27D + 81E = 10, \\ A + 4B + 16C + 64D + 256E = 20, \\ A + 5B + 25C + 125D + 625E = 35. \end{cases}$$

From these equations we find the values of A, B, C, D, E ; whence,

$$U_n = -20 + 36n - \frac{115n^2}{6} + \frac{9n^3}{2} - \frac{n^4}{3}.$$

From this we get the 6th term to be 46.

Thus, the series is

$$1 + 6 + 10 + 20 + 35 + 46 + \dots$$

Using the method of differences we get

$$\begin{array}{ccccccc} 5 & 4 & 10 & 15 & 11 & \dots \\ -1 & 6 & 5 & -4 & \dots & \\ 7 & -1 & -9 & \dots & \dots & \\ -8 & -8 & \dots & \dots & \dots & \\ 0 & \dots & \dots & \dots & \dots & \end{array}$$

Then

$$\begin{aligned} S_n &= n + \frac{5n(n-1)}{|2|} - \frac{n(n-1)(n-2)}{|3|} + \frac{7n(n-1)(n-2)(n-3)}{|4|} - \frac{8n(n-1)(n-2)(n-3)(n-4)}{|5|} \\ &= -\frac{n}{120} (8n^4 - 115n^3 + 510n^2 - 1145n + 622). \end{aligned}$$

(c) In the series $1 + 5 + 15 + 35 + 70 + \dots$

$$U_n = A + Bn + Cn^2 + Dn^3 + En^4 + \dots$$

$$\left\{ \begin{array}{l} A + B + C + D + E = 1, \\ A + 2B + 4C + 8D + 16E = 5, \\ A + 3B + 9C + 27D + 81E = 15, \\ A + 4B + 16C + 64D + 256E = 35, \\ A + 5B + 25C + 125D + 625E = 70, \end{array} \right\} \begin{array}{l} A = 0, \\ B = 1/4, \\ C = 11/24, \\ D = 1/4, \\ E = 1/24. \end{array}$$

Hence,

$$U_n = \frac{n}{4} + \frac{11n^2}{24} + \frac{n^3}{4} + \frac{n^4}{24}.$$

Then

$$24S_n = 6\Sigma n + 11\Sigma n^2 + 6\Sigma n^3 + \Sigma n^4,$$

$$24S_n = 3n(n+1) + \frac{11n(n+1)(2n+1)}{6} + \frac{3n^2(n+1)^2}{2} + \frac{n(n+1)(6n^3+9n^2+n-1)}{30}.$$

$$S_n = \frac{n}{120} (n^4 + 10n^3 + 35n^2 + 50n + 24) = \frac{n}{120} (n+1)(n+2)(n+3)(n+4).$$

Also solved by O. S. ADAMS, H. H. CONWELL, PAUL CAPRON, and WILLIAM TIER.

GEOMETRY.

496. Proposed by NATHAN ALTSHILLER, University of Oklahoma.

Find all the lines such that the pairs of tangent planes to a given sphere (ellipsoid) passing through them, shall be orthogonal.

SOLUTION BY WILLIAM HOOVER, Columbus, Ohio.

One obvious solution is given when one of the tangent planes is fixed in position, for then this plane is the locus of lines common to it and a second tangent plane orthogonal to the first.

It is not troublesome to show that the equation of a first plane embracing a line

$$\frac{x-a}{l} = \frac{y-b}{m} = \frac{z-c}{n}, \quad (1)$$

$$\frac{(x-a)\lambda_1}{l} + \frac{(y-b)\mu_1}{m} + \frac{(z-c)\nu_1}{n} = 0, \quad (2)$$